

Research Article

# Peculiarities of the Course and Provision of Medical Care to Patients with Acute Coronary Syndrome in the Precarpathian Region

Roksolana Nesterak

## Abstract

**Abstract.** Cardiovascular diseases are the first ones in mortality among other diseases. Particular attention should be paid to patients after acute coronary syndrome, since their quality of life, the possibility of restoration of work capacity.

**Objective of the research.** To evaluate indicators of medical care provision for patients with acute coronary syndrome in the Precarpathian region.

**Materials and methods.** The medical-geographical indicators, data of the local registry (2014-2018) included clinical and objective parameters, peculiarities of the course, rehabilitation and treatment of patients with acute coronary syndrome.

**Results.** The peculiarities of the course of acute coronary syndrome, clinical characteristics of patients depend on the form of IHD and the applied method of ACS treatment. There is an increase of the number of PCI performance and the reduction of TLT, the percentage of admission of patients up to 2 hours remains low. The increase of PCI performed in the districts of the region in the dynamics from 2014 to 2018 has been determined, as well as the geographical peculiarities of the Ivano-Frankivsk region. The percentage of patients' undergoing rehabilitation after acute coronary syndrome remains low.

**Conclusions.** The Precarpathian region has a number peculiarities that are related to the location of the region. Taking into account the characteristics will contribute to the improvement of the organizational model of medical care provision in the region.

## Keywords

acute coronary syndrome; Precarpathian Region

*Ivano-Frankivsk National Medical University, Ivano-Frankivsk, Ukraine*

**Corresponding author:** roxolana.nesterak@gmail.com



Copyright ©Roksolana Nesterak, 2019

## Problem statement and analysis of the latest research

The ischemic heart disease is the most frequent reason of death all over the world. Over seven million people with IHD die every year, that is 12.8% out of all mortal cases [1].

The frequency of cases of hospitalization with myocardial infarction with the ST elevation is various in different countries – the participants of the European Society of Cardiologists. NSTEMI in the


long dates has mortality indices similar to STEMI, which causes attention to these patients in the acute phase, and in the period of rehabilitation [2, 3].

For the valuation of the adherence to recommendations and the effectiveness of treatment of patients in different countries the registers are used – special scientific research programmes, in which all patients with pathology being studied, are included [4, 5, 6]. During the last years such registers are one of the main ways of studying of acute coronary



of disability) and life (more than half of all cases of death) are circulatory system diseases (CSD): atherosclerosis, ischemic heart disease (IHD), arterial hypertension (AH) and their complications (heart attack, stroke), which rank first among other nosologies.

To the structure of total mortality of the population, in which during 2000-2018 growth for diseases of the digestive system (+ 40.0%), cholesterol (+19.5%), onco- (+4.9%) and decrease in respiratory diseases (-82.0%) was noted (Fig. 1).



**Figure 1.** Health indicators of the Ivano-Frankivsk region during 2000-2018.

At the age of 45-59, the main percentage is men 27 (27.5%), women 4 - (4.08%) with NSTEMI and 65 (23.5%) and 9 (3.2%) at STEMI, respectively; in the age group 60-74, the proportion of women with NSTEMI 16 (16.3%) and STEMI 52 (16.84%) versus men is 20 (20.4%) and 16 (16.33%), and at the age above 75 years for NSTEMI women prevail - 16


(16.3%). This trend was observed in previous years and is comparable to the general data of Ukraine. The distribution of patients with NSTEMI by gender and age is represented in Figure 2, STEMI (Figure 3).

Analyzing the clinical characteristics of patients with acute coronary syndrome, it was found that among patients with NSTEMI, which used conservative treatment, 63.3% (38) were residents of the city, 36.7% - residents of the village; respectively, the PCI was performed in 68.0% and 32.0% of patients (Table 1). In a cohort of patients - residents of the city with STEMI 55, 6% of patients treated conservatively, in one third of patients (35, 1%) – thrombolysis was performed and in almost half of patients (46, 2%) – PCI was performed. Accordingly, the villagers accounted for 4.4%, 66.7% and 53.9%, namely, 2/3 of patients who used TLT were residents of the village.


The percentage of patients without increase in blood pressure was insignificant, and it was in the group of patients with STEMI 9.5%, 10.5% and 10.2%. In most of the patients examined there was an arterial hypertension (AH). Moreover, in spite of the applied medical technology and the course of coronary artery disease, the majority of patients had AH II. In the presence of NSTEMI, the 3<sup>rd</sup> degree of hypertension was observed in 21.7% and 25.3% of patients with conservative treatment and PCI. TLT in the case of STEMI was used in 10.5% of patients without hypertension and in 8.8%, 68.4% and 12.3% of patients with hypertension of the first to third degree, respectively. In all groups, patients were associated with diabetes type II, the frequency of which was – from 16.0% to 31.7% of cases

In the region between 2014 and 2018 there is a decrease in cases of myocardial infarction (Fig. 4). So, the highest level was in 2016 – 1297 cases, in 2018 – 1188 cases. No significant fluctuations in percentages are observed for STEMI and NSTEMI cases. STEMI in 2016 was 962 cases (74.1%), and in 2018 – 815 cases (68.3%). NSTEMI in 2016 – 336 cases (25.9%), and in 2018 – 373 cases (31.7%).

From 2015, there is an increase in the percentage of patients who have been diagnosed with PCI.



**Figure 2.** Distribution of patients with NSTEMI by age and gender.



**Figure 3.** Distribution of patients with STEMI by age and gender.

**Table 1.** The clinical characteristics of patients with acute coronary syndrome.

	NSTEMI (n=135)		STEMI (n=198)		
	Conservative treatment (n=60)	PCI (n=75)	Conservative treatment (n=63)	TLT (n=57)	PCI (n=78)
Residents of the city	38 (63.33%)	51 (68.00%)	35 (55.56%)	20 (35.09%)	36 (46.15%)
Residents of the village	22 (36.67%)	24 (32.00%)	28 (44.44%)	37 (66.67%)	42 (53.85%)
Without AH	9 (15.00%)	5 (6.67%)	6 (9.52%)	6 (10.53%)	8 (10.26%)
AH I	-	3 (4.00%)	-	5 (8.77%)	2 (2.56%)
AH II	38 (63.33%)	48 (64.00%)	43 (68.25%)	39 (68.42%)	43 (55.13%)
AH III	13 (21.67%)	19 (25.30%)	14 (22.22%)	7 (12.28%)	25 (32.05%)
Diabetes type II	19 (31.67%)	12 (16.00%)	11 (17.46%)	10 (17.54%)	15 (19.23%)
Without heart failure			4 (6.35%)	5 (8.77%)	15 (19.23%)
HF I	51 (85.00%)	54 (72.00%)	9 (14.29%)	8 (14.04%)	7 (8.97%)
HF IIA	8 (13.33%)	14 (18.67%)	18 (28.57%)	17 (29.82%)	17 (21.79%)

*Notes:*

1. The absolute number of patients is indicated.
2. The percentage to the absolute number of studied persons is represented in the brackets.

So, in 2015, 333 patients with ACS were followed by PCI, in 2018 there were 582 interventions. Due to the geographical features and remoteness of some districts from the only one center in the region where primary PCI is performed, in part of patients, thrombolytic therapy (TLT) was used. There were 75 TLT in 2018 against 175 in 2014 (Fig. 5).

Taking into account the geographical features of the region, as well as the insufficient number of centers with the possibility of PCI performing, 343 patients with ACS in 2018 were admitted to the united center of Ivano-Frankivsk region: up to 2 hours – 66 patients – 21%, 2-6 hours – 162 patients – 51%, 7-24 hours of patients – 19 – 6%, more than

24 hours – 19 patients – 6% (Fig. 6).

Analyzing the amount of the performed PCI in the districts of the region, the increase of the number of the intervention was defined from 2014 till 2018 (Fig. 7). However, the features of geographical position and the great distance of the districts from the PCI, centres with the possibility of providing had been marked. So, the smallest number of the patients, who were given PCI, live in Verhovynsky district – 3 (three) and in the town of Bolekhiv – 4 (four). The largest number, given PCI was in 2018 in the city of Ivano-Frankivsk – 187 (one hundred and eighty-seven).

With the purpose of improvement of giving med-

Peculiarities of the Course and Provision of Medical Care to Patients with Acute Coronary Syndrome in the Precarpathian Region — 6/10




Figure 4. The number of acute myocardial infarction 2014-2018.





Figure 5. Dynamics of ACS treatment.



**Figure 6.** Structure of patients by time of admission to the center with possibility of PCI.

ical aid it is expedient to establish the centres with conducting invasive diagnostic methods, treatment and rehabilitation of the patients in the following towns. In particular, in the city of Ivano-Frankivsk, including Ivano-Frankivsk and the following districts:

- Halytsky, Tlumatsky, Tysmenetsky, Rohatynsky and the town of Burshtyn;
- in the town of Kalush, including such districts: Kalusky, Rozhniativsky, Dolynsky and the town of Bolekhiv;
- in the town of Kolomyia, including the following districts: Kolomyysky, Horodenkivsky and Sniatynsky;
- in the town of Nadvirna, including such districts: Nadvirniansky, the town of Yaremche and Bohorodchansky district.

Kosivsky centre can be separately allocated. It will unite Kosivsky and Verhovynsky districts.

In particular, in Verhovynsky district – the villages of Holoshyna (54km), Burkut (37km), in Rohatynsky district – the villages Melna (25km), Zhuravensky (33km), Svitanok (37km), in Dolynsky district – the villages of Vyshkiv (43km), Senechiv (47km) and village of Trostianets (32km) in Sniatynsky district.

The rehabilitation and the restoration is the inte-

gral part of supplying the patients with acute coronary syndrome with effective medical care.

On the basis of the Ivano-Frankivsk Regional Clinical Cardiology Center there is a rehabilitation department where patients of working age underwent rehabilitation after acute coronary syndrome and cardiac surgery.

In the region there is a need for a rehabilitation institution, where provision of more effective rehabilitation for the entire population of the region. The percentage of rehabilitation is low, for example in 2017 – 28.1% of patients (Fig. 8).

There is a multidisciplinary approach in the rehabilitation department, which covers all components of rehabilitation. Physical rehabilitation is done (including morning gymnastics, medical physical exercises, closed walking, physiotherapy, procedures, massage) along with medical aid.

The suggestive methods (auto-training, music therapy, individual and group counseling, training are used in psychological rehabilitation. The teaching of the patients is one of the methods used in the department. A school-club "Health Heart" was created. To support patients in the medical and psychological areas; increase of awareness of patients with cardiovascular diseases psychological component of diseases; improve of rehabilitation of




Figure 7. The amount of PCI from 2014 till 2018 in the district of Ivano-Frankivsk region.




Besides taking into account the remoteness of the districts from the regional centre or the centre with the possibility to conduct PCI, it's necessary to take into account the most remote villages in each district.

patients; increase patients' attachment to non-drug and medical treatment; work on the internal picture of health and the internal picture of the disease; form skills and abilities of influence on behavioural risk factors; train self-control skills; provide the first premedical care; promote healthy lifestyle, increasing the patient's responsibility for maintaining his health and motivating him to recover; work with families and relatives of patients is the purpose of the work. A page of social networks is also created for social awareness.

### 3. Conclusions

1. The cardiovascular diseases take the first place in mortality among other diseases despite the introduction of new diagnostic methods and treatment. The acute coronary syndrome, as the manifestation of ischemic heart disease, leads to disability, reducing the quality and life expectancy of patients. It needs improvement of the methods of treatment, in particular the rehabilitation and restorative treatment



**Figure 8.** Rehabilitation after the acute coronary syndrome in the districts.

of such patients.

2. Taking into account the importance of studying and improvement of the quality of provision of medical care for the patients with the cardiovascular diseases it is relevant to take into account the geographical situation of the districts of Ivano-Frankivsk region.
3. The geographical peculiarities, reducing attention of the population to the health, appeal for medical treatment in critical situations, failure to conduct the preventive or restorative treatment – all this is the peculiarity of the Precarpathian region.
4. The determined regional peculiarities and the remoteness of the districts of Ivano-Frankivsk region from the regional, centre, when PCI and the surgical revascularization are conducted, it is necessary to create the centres with the possibility to conduct the revascularization.

#### 4. Prospects of Further Researches

Perspective is the peculiarities of the course and provision of medical care to patients with stable ischemic heart disease in the Precarpathian region.

#### References

- [1] Unified clinical protocol of emergency, primary, secondary, third medical care and medical rehabilitation "Acute coronary syndrome without elevation of segment ST", Order of the Ministry of Health of Ukraine 03.03.2016 #164.
- [2] Aglullina EI. Acute coronary syndrome: from diagnosis to optimization of treatment. *Vestnik sovremennoy klinicheskoy meditsyny*. 2013; 6 (5): 91-94. DOI: [https://doi.org/10.20969/VSKM.2013.6\(5\).91-94](https://doi.org/10.20969/VSKM.2013.6(5).91-94)
- [3] Terenda NO, Petrashyk YuM. Comparative assessment of the effectiveness of the interven-

tional method of treating patients with myocardial infarction according to the Ternopil region. Hospital Surgery. Journal named after LYa. Kovalchuk 2017; 1: 91-95.

- [4] Ferreira J, Monteiro P, Mimoso J. National Registry of Acute Coronary Syndromes: results of the hospital phase in 2002. *Rev. Port. Cardiol.* 2004; 23 (10):1251-1272.
- [5] Pajunen P, Koukkunen H, Ketonen M. et al. The validity of the Finnish Hospital Discharge Register and Causes of Death Register data on coronary heart disease. *Eur. J. Cardiovasc. Prev. Rehabil.* 2005;12: 132-137. DOI: <https://doi.org/10.1097/00149831-200504000-00007>
- [6] Garganeeva AA, Okrugin CA, Borel KN. The WHO Register of Acute Myocardial Infarction is an opportunity and perspective in the study and forecasting outcomes of socially significant poatologies at the population level. *Sibirskiy meditsinskiy zhurnal.* 2015; 2: 125-130.
- [7] Efros LA, Samorodskaya IV. Survival and efficiency in men after coronary bypass. *Clinical medicine.* 2013; 5: 27-31.
- [8] Azizov VB, Antonyuk IV, Baranenko AV, et al. Registry of Percutaneous Coronary Intervention: Comparative Analysis 2014-2015 Regional reperfusion networks in Ukraine - the dynamics of development. *Sertse i sudyny.* 2016; 3: 14-34.
- [9] Kovalenko VN, Dolzhenko MN, Nesukai EG, Dyachenko Ya. Comparative characteristics of prevention of cardiovascular diseases in Ukraine and Europe according to EUROASPIRE VI: hospital line. Arterial hypertension. 2016; 1 (45): 57-61. DOI: <https://doi.org/10.22141/2224-1485.1.45.2016.74138>
- [10] Chendey TV, Ravtochka YaH, Lohoyda VV. Register of acute coronary syndromes in Transcarpathia: first results. *Sertse i sudyny.* 2017; 2: 38-44.
- [11] Geographical location and natural conditions of the Carpathian region. Ivano-Frankivsk Regional Universal Library named after I. Franko. <https://lib.if.ua/exhib/1534076504.html>
- [12] Yatsuba VG, et al. Administrative-territorial structure of Ukraine. History and the present. *KIYV.* 2001; 272.

**Received:** 2019-05-13

**Revised:** 2019-05-24

**Accepted:** 2019-06-02