Short Report

Intermammary Pilonidal Disease

Arda Isik*, Eray Kurnaz, Nurten Arslan Isik

Erzincan University, Turkey

*Corresponding author: kararda@yahoo.com

Copyright ©Arda Isik, Eray Kurnaz, Nurten Arslan Isik, 2019

Pilonidal disease is an abnormal skin growth that contains hairs. It occurs mainly in the sacro-coccygeal area; however, it may rarely occur in the axilla, groin, interdigital web space or umbilicus. In the literature, there were only 13 cases pilonidal disease in the intermammary region [1, 2].

Here we present a case of a 22-year-old female. She had chronic discharging sinus in the intermammary area for six month. The examination revealed that the patient was hirsute and had multiple sinuses in the intermmarian region (Fig. 1).

Serous discharge was present in the patient; however, there was no abcess formation. The patient was treated with antibiotics and curettage of the sinus epithelium. After one week, the discharge stopped. During the 6-month follow-up, there was no recurrence.

Figure 1. Multiple sinuses in the intermmarian region.

References

- [1] Seerwan Hama Shareef, Tahir A Hawrami, Abdulwahid M Salih et al. Intermammary pilonidal sinus: The first case series. *Int J Surg Case Rep.* 2017;41:265-268. DOI: <https://doi.org/10.1016/j.ijscr.2017.10.021> [PMid:29121582 PMCID:PMC5681334]
- [2] Anil Sunkara, DD Wagh, Sameer Harode. Intermammary Pilonidal Sinus. *Int J Trichology.* 2010;2:116-118. DOI: <https://doi.org/10.4103/0974-7753.77526> [PMid:21712902 PMCID:PMC3107957]

Received: 2019-05-21

Revised: 2019-06-26

Accepted: 2019-06-26